

Surface Geology

North Killdeer Mountain Quadrangle, North Dakota

Edward C. Murphy
2007

EXPLANATION

QUATERNARY SYSTEM

RECENT

OAHE FORMATION

Qal Alluvium

Brownish gray to black sand, silt, clay, and lenses of gravel; flood-plain deposits along recent drainages. Includes lower terrace deposits. Typically less than 50 feet thick.

RECENT/PLEISTOCENE

Qls Landslide Deposits

Variable mixture of strata and deposits that have slid to the base of steep slopes.

PLEISTOCENE

COLEHARBOR GROUP

Qp Pediment Deposits

Variable mixture of strata and deposits that have slid to the base of steep slopes.

TERTIARY SYSTEM

MIOCENE/OLIGOCENE

Ta ARIKAREE FORMATION

Buff-colored tuffaceous sandstone, siltstone, and fresh water carbonate.

EOCENE

Tc CHADRON FORMATION

Sandy and pebbly mudstone and clayey sandstone.

EOCENE/PALEOCENE

Tgv GOLDEN VALLEY FORMATION

Camels Butte Member

Alternating beds of yellowish brown to brown, micaceous sandstone, siltstone, mudstone, claystone, and lignite.

Bear Den Member

Brightly colored, kaolinitic claystone, mudstone, and sandstone, typically overlain by a thin siliceous bed (silerete) or lignite.

PALEOCENE



Tsb SENTINEL BUTTE FORMATION





Alternating beds of grayish brown to gray sandstone, siltstone, mudstone, claystone, and lignite.

Geologic Symbols

- Known contact between two geologic units.
- - - Approximate contact between two geologic units.
- - - - Inferred (unknown) contact between two geologic units.

Other Features

-  Water
-  Unpaved Road



Lambert Conformal Conic Projection Standard Parallels 47° 30' 00" and 47° 37' 30"
1927 North American Datum NGVD 1929
USGS 7.5 Minute Topographic Map Contour Interval 20 Feet
Road Layer Recultured to 2003 NAIP Digital Orthophoto

